

UNIONS OF CUBES IN $\mathbb{R}^n$, COMBINATORICS IN $\mathbb{Z}^n$ AND THE JOHN-NIRENBERG AND JOHN-STRÖMBERG INEQUALITIES

MICHAEL CWIKEL

ABSTRACT. Suppose that the d -dimensional unit cube Q is the union of three disjoint “simple” sets E , F and G and that the volumes of E and F are both greater than half the volume of G . Does this imply that, for some cube W contained in Q , the volumes of $E \cap W$ and $F \cap W$ both exceed s times the volume of W for some absolute positive constant s ?

Here, by “simple” we mean a set which is a union of finitely many dyadic cubes.

We prove that an affirmative answer to this question would have deep consequences for the important space BMO of functions of bounded mean oscillation introduced by John and Nirenberg.

The notion of a John-Strömberg pair is closely related to the above question, and the above mentioned result is obtained as a consequence of a general result about these pairs. We also present a number of additional results about these pairs.

1. INTRODUCTION

One of the aims of this paper is to warmly invite readers with interests quite unrelated to the particular realm in analysis which gave rise to the geometrical and almost combinatorial question being considered here, to study that question, and hopefully even resolve it. In particular (as our title seeks to indicate, and as confirmed by [4]) those readers with expertise in geometry in $\mathbb{R}^n$ or combinatorial problems in $\mathbb{Z}^n$ may well have some valuable insights. Those who choose to respond to this invitation will be able to effectively do so, without needing any familiarity whatsoever with the proofs in this paper, or with the contents of the earlier papers which gave rise to it.

This paper is a sequel to [3] and its preliminary more detailed version [1]. The main motivation for those papers, and also for this one, is the wish for a better understanding of the celebrated John-Nirenberg inequality [5] which is satisfied by functions $f : D \rightarrow \mathbb{R}$ in the class $BMO(D)$ of bounded mean oscillation defined on a suitable subset D of $\mathbb{R}^d$. In particular, it is hoped to ultimately find an answer to:

Question J-N. *Can the constants in the John-Nirenberg inequality for BMO functions of d variables be chosen to be independent of d ?*

We refer to [6], [7], [8] and [9], and also to references in these papers. for some recent results concerning best constants in certain versions of the John-Nirenberg inequality, mainly in the case where $d = 1$. Some remarks on pp. 7–8 of [7] recall some reasons

Key words and phrases. BMO , John-Strömberg pairs, John-Nirenberg inequality.

This research was supported by funding from the Martin and Sima Jelin Chair in Mathematics, by the Technion V.P.R. Fund and by the Fund for Promotion of Research at the Technion.

for being interested in the sizes of these constants, also for $d > 1$. We are grateful to Andrei Lerner and Leonid Slavin for information about these and related results.

We shall present a new geometrical, almost combinatorical question which is quite easy to formulate, expressed in terms readily accessible to a general mathematical audience, and whose affirmative answer would imply an affirmative answer to Question J-N. We will refer to it as Question A(1/2). It would seem to be at least slightly easier to answer than certain very similar questions which have an analogous role and which are posed in [1] and [3], and are also discussed in the brief survey article [2]. In the formulation of Question A(1/2), and indeed throughout this paper, we shall use the following notation and terminology (much of which is standard, and most of which was also used in [1] and [3]).

Definition 1.1. We shall understand that

- (i) d is a positive integer, and that
- (ii) the word **cube** means a closed cube in $\mathbb{R}^d$ with sides parallel to the axes, i.e., the cartesian product of d bounded closed intervals, all of the same length. Special roles will be played by the d -dimensional **unit cube** $[0, 1]^d$ and its **dyadic subcubes** (i.e, the cubes of the form $\prod_{j=1}^d [(n_j - 1)2^{-k}, n_j 2^{-k}]$, where $k \in \mathbb{N}$ and the integers n_j all satisfy $1 \leq n_j \leq 2^k$). It will sometimes be convenient to use the standard notation $Q(x, r)$ for the cube of side length $2r$ centred at $x \in \mathbb{R}^d$, i.e., $Q(x, r) = \{y \in \mathbb{R}^d : \|y - x\|_{\ell_d^\infty} \leq r\}$.

Furthermore,

- (iii) we let $\mathcal{Q}(\mathbb{R}^d)$ denote the collection of all cubes in $\mathbb{R}^d$, and
- (iv) the word **d -multi-cube** will mean a non-empty subset of $[0, 1]^d$ which is the union of finitely many **dyadic subcubes** of $[0, 1]^d$, and
- (v) we will denote the d -dimensional Lebesgue measure of any Lebesgue measurable subset E of $\mathbb{R}^d$ by $\lambda(E)$.

(The values of d in (ii) and in (v) will be clear from the contexts in which they appear.)

In fact, in the formulation of Question A(1/2) we will not need any subtle properties of Lebesgue measure. The only Lebesgue measurable subsets E of $\mathbb{R}^d$ that we will encounter in it are cubes or d -multi-cubes and their intersections with other cubes. All of these are finite unions $E = \bigcup_{m=1}^M E_m$ of non-overlapping sets E_m , where each E_m is the cartesian product of d closed intervals. So we simply have that each $\lambda(E_m)$ is the product of the lengths of those d intervals, and that $\lambda(E) = \sum_{m=1}^M \lambda(E_m)$. With some small adjustments, we can replace consideration of $\lambda(E)$ for all the above mentioned sets E by consideration of the cardinality of $E \cap 2^{-k}\mathbb{Z}^d$ for a suitably large choice of $k \in \mathbb{N}$. (This possibility was already very briefly hinted at in [2].) This is one reason for our suggestion above, in the opening paragraph of this paper, that Question A(1/2) might perhaps also be approachable via combinatorical considerations.

After these preparations, here at last is the promised question:

Question A(1/2). *Does there exist an absolute constant $s > 0$ which has the following property?*

For every positive integer d , whenever E_+ and E_- are two disjoint d -multi-cubes which satisfy

$$(1.1) \quad \min \{ \lambda(E_+), \lambda(E_-) \} > \frac{1}{2} (1 - \lambda(E_+) - \lambda(E_-)) ,$$

then there exists a cube W which is contained in $[0, 1]^d$ and for which

$$(1.2) \quad \min \{ \lambda(W \cap E_+), \lambda(W \cap E_-) \} \geq s \lambda(W) .$$

Most of the work which is required to show that an affirmative answer to Question A(1/2) implies an affirmative answer to Question J-N has already been done in [1] and [3]. Because of that we will not have any need at all here to deal with any details concerning any of the versions of the function space BMO , nor to even recall their definitions. It was shown in [1] and [3] that an affirmative answer to Question J-N would be a consequence of an affirmative answer to the following question which was formulated at the beginnings of both of those papers, and is clearly quite similar to Question A(1/2) above.

Question A. *Do there exist two absolute constants $\tau \in (0, 1/2)$ and $s > 0$ which have the following property?*

For every positive integer d and for every cube Q in $\mathbb{R}^d$, whenever E_+ and E_- are two disjoint measurable subsets of Q whose d -dimensional Lebesgue measures satisfy

$$\min \{ \lambda(E_+), \lambda(E_-) \} > \tau \lambda(Q \setminus E_+ \setminus E_-) ,$$

then there exists a cube W which is contained in Q and for which

$$\min \{ \lambda(W \cap E_+), \lambda(W \cap E_-) \} \geq s \lambda(W) .$$

It is obvious that an affirmative answer to Question A would imply an affirmative answer to Question A(1/2). Our task here is to show that the reverse implication also holds, namely that affirmatively answering the apparently at least slightly less demanding Question A(1/2) would suffice to also affirmatively answer Question A and therefore also Question J-N.

After recalling some more notions and providing some preliminary results in Section 2, we will obtain the above mentioned reverse implication in Section 3, as an immediate consequence (see Corollary 3.2 below) of the main result, Theorem 3.1 of this paper, which is set in a slightly more general context than Questions A and A(1/2).

We note that the result of Theorem 3.1 could also conceivably make it possible to deduce an affirmative answer for Question J-N from certain variants of Question A, which might perhaps be easier to answer than Question A(1/2). Some explorations of the possibility of such options, including some relevant numerical experiments, will perhaps be discussed in a future update of this paper.

The above mentioned more general context in which Theorem 3.1 is formulated revolves around the notion of John-Strömberg pairs. In Section 4 we shall offer a number of additional results about these pairs, beyond those which will be needed for our main result.

Finally, the extremely brief Section 5 offers some comments about the papers [1] and [3].

Remark 1.2. It is a pleasure to describe three results which were recently obtained by Ron Holzman [4]. The first of these is an affirmative answer to a non-trivial special case of Question A(1/2): Holzman has shown, for every $d \in \mathbb{N}$, that whenever E_+ and E_- are d -multi-cubes which are each finite unions of dyadic cubes, *all of side length* $1/2$, and they satisfy (1.1), then there exists a cube W in $[0, 1]^d$ which satisfies (1.2) for $s = 1/4$. In fact, as will be explained below in Remark 4.7, this value of s is apparently “optimal” in the sense that, if the general form of Question A(1/2) has an affirmative answer, then the positive number s which appears in that answer must satisfy $s \leq 1/4$.

The second of Holzman’s recent results is that this largest possible value $1/4$ for s is indeed attained in another special case of Question A(1/2), where d is restricted to take only one value, namely $d = 1$, but where there is no restriction on the form of the disjoint measurable subsets E_+ and E_- of $[0, 1]$. (See also Remark 4.10.) Holzman’s third result (see Remark 4.7 for details) shows that the analogue of his second result, when the single chosen value for d is greater than 1, does not hold. For each $d \geq 2$, the relevant value of s cannot be greater than $\sqrt{5} - 2$. Thus this smaller number is now revealed as a better upper bound for the positive number s , should there be an affirmative answer for Question A(1/2) for that s .

2. SOME FURTHER DEFINITIONS AND PRELIMINARY RESULTS

The following definition (which is effectively the same as Definition 7.10 of [1, p. 29]) recalls a notion which plays a central role in [1] and [3]. and which is of course closely related to Question A.

Definition 2.1. Let d be a positive integer and let $\mathcal{E}$ be a non-empty collection of Lebesgue measurable subsets of $\mathbb{R}^d$. Suppose that each $E \in \mathcal{E}$ satisfies $0 < \lambda(E) < \infty$. Let τ and s be positive numbers with the following property:

(*) *Let Q be an arbitrary set in $\mathcal{E}$ and let E_+ and E_- be arbitrary disjoint measurable subsets of Q . Suppose that*

$$(2.1) \quad \min \{ \lambda(E_+), \lambda(E_-) \} > \tau \lambda(Q \setminus E_+ \setminus E_-).$$

Then there exists a set $W \subset Q$ which is also in $\mathcal{E}$ and for which

$$(2.2) \quad \min \{ \lambda(E_+ \cap W), \lambda(E_- \cap W) \} \geq s \lambda(W).$$

Then we will say that (τ, s) is a **John-Strömberg pair** for $\mathcal{E}$.

Remark 2.2. In the context of the preceding definition, if E_+ , E_- and W are measurable subsets of Q which satisfy $E_+ \cap E_- = \emptyset$ and $\lambda(W) > 0$ and (2.2) for some $s > 0$, then

$$\lambda(W) = \lambda(E_+ \cap W) + \lambda(E_- \cap W) \geq 2 \min \{ \lambda(E_+ \cap W), \lambda(E_- \cap W) \} \geq 2s \lambda(W) > 0$$

and therefore $s \leq 1/2$. Consequently, for any choice of $\mathcal{E}$, any pair (τ, s) which is a John-Strömberg pair for $\mathcal{E}$ must satisfy $0 < s \leq 1/2$. (Cf. Remark 7.16 of [1, p. 30].)

Several “natural” choices of the collection $\mathcal{E}$ are mentioned on pp. 4–5 of [1] and pp. 132–133 of [3]. These are relevant for studying a number of variants of the function

space BMO which have been considered in the literature. The most “classical” of these choices, motivated by the context of [5] and by Question J-N, is when we take $\mathcal{E}$ to be the collection $\mathcal{Q}(\mathbb{R}^d)$ of all cubes in $\mathbb{R}^d$ (as introduced in Definition 1.1(iii)) for some fixed $d \in \mathbb{N}$.

It is known that $(\sqrt{2} - 1, 2^{-d}(3 - 2\sqrt{2}))$ is a John-Strömberg pair for $\mathcal{Q}(\mathbb{R}^d)$. (See Example 7.12(i) on p. 30 of [1].) Indeed in this paper we will need to use Definition 2.1 only in the special case where $\mathcal{E}$ is the collection $\mathcal{Q}(\mathbb{R}^d)$. But we have indicated the possibility of considering other choices of the collection $\mathcal{E}$, since there may turn out to be variants of our main result in this paper, and these may turn out to be relevant for future research in the contexts of at least some of those other choices. (We refer to [1] and [3] for various results, in particular Theorem 9.1 of [1, p. 41] and [3, p. 164], which deal with John-Strömberg pairs and their interactions with versions of the John-Nirenberg inequality for variants of the space BMO corresponding to other choices of $\mathcal{E}$.)

Definition 2.3. It will sometimes be convenient, for each $d \in \mathbb{N}$, to let $JS(d)$ denote the set of all ordered pairs (τ, s) of positive real numbers which are John-Strömberg pairs for $\mathcal{Q}(\mathbb{R}^d)$.

Obviously, an equivalent reformulation of Question A is:

Question A (τ, s) . *Do there exist two absolute constants $\tau \in (0, 1/2)$ and $s > 0$ for which $(\tau, s) \in JS(d)$ for every $d \in \mathbb{N}$?*

This makes it very relevant to use the following known result. (Note that below in Subsection 4.1 we will present a result containing some new slight variants of it.)

Theorem 2.4. *Let d be a positive integer. The ordered pair of positive numbers (τ, s) is a John-Strömberg pair for $\mathcal{Q}(\mathbb{R}^d)$ if and only if it has the following property:*

(*) *Whenever F_+ and F_- are disjoint d -multi-cubes which satisfy*

$$(2.3) \quad \min \{ \lambda(F_+), \lambda(F_-) \} > \tau \lambda([0, 1]^d \setminus F_+ \setminus F_-),$$

then there exists a cube W contained in $[0, 1]^d$ for which

$$(2.4) \quad \min \{ \lambda(W \cap F_+), \lambda(W \cap F_-) \} \geq s \lambda(W).$$

Proof. Obviously every John-Strömberg pair (τ, s) for $\mathcal{Q}(\mathbb{R}^d)$ must have the property (*). The fact that property (*) implies that (τ, s) is a John-Strömberg pair for $\mathcal{Q}(\mathbb{R}^d)$ is exactly the content of Theorem 10.2 of [1, p. 43], and is proved on pp. 44–47 of [1]. However, some other small issues remain to be clarified: It should be mentioned that there are four minor misprints in the proof of Theorem 10.2 of [1]. But they do not effect its validity. On page 46, in two places (in the third line and in the fourth line above the inequality (10.16)) the notation F_ε should of course be F_b . Then on line 11 of page 47 r_n^d should be $(2r_n)^d$ and r_*^d should be $(2r_*)^d$. Similarly, four factors of 2^d have been (harmlessly) omitted on line 5 of page 63 of [1] in the proof of Lemma 10.1, which is an ingredient in the proof of Theorem 10.2. That line should be

$$= (2R_n)^d - (2r_*)^d + (2R_n)^d - (2r_n)^d.$$

It should also be mentioned that, in the formulation of Theorem 10.2 of [1], it is stated that the number τ must satisfy $0 < \tau < 1/2$. But the proof is valid for all $\tau > 0$. (The

requirement that $\tau < 1/2$ was imposed only because this is relevant in Theorem 9.1 of [1, p. 41] and [3, p. 164].)

The clarification of these small issues completes the proof of Theorem 2.4. $\square$

3. THE MAIN RESULT

We can now present our main result, and its obvious corollary for dealing with Question A(1/2)

Theorem 3.1. *Let d be a positive integer. Suppose that (τ, s) is a John-Strömberg pair for $\mathcal{Q}(\mathbb{R}^d)$. Then, for each $\theta \in (0, s/(1-s))$, the pair $((1-\theta)\tau, s-\theta(1-s))$ is also a John-Strömberg pair for $\mathcal{Q}(\mathbb{R}^d)$.*

Corollary 3.2. *Suppose that the answer to Question A(1/2) is affirmative, i.e., suppose that $(1/2, s)$ is a John-Strömberg pair for $\mathcal{Q}(\mathbb{R}^d)$ for some positive number s which does not depend on d . Then, if we choose $\theta = s/(2-2s)$ in Theorem 3.1, we obtain that $(\frac{2-3s}{4-4s}, \frac{s}{2})$ is a John-Strömberg pair for $\mathcal{Q}(\mathbb{R}^d)$ for all $d \in \mathbb{N}$, which gives us an affirmative answer to Question A(τ, s) and therefore also to Question A and Question J-N.*

Proof of Theorem 3.1. Let us explicitly choose particular (necessarily positive) numbers τ , s and θ which satisfy the hypotheses of the theorem.

As observed in Remark 2.2, the fact that (τ, s) is a John-Strömberg pair for $\mathcal{Q}(\mathbb{R}^d)$ ensures that $0 < s \leq 1/2$. So the conditions imposed on θ ensure that $0 < \theta < 1$ and also that $\theta(1-s) < s$. Consequently, the numbers $(1-\theta)\tau$ and $s-\theta(1-s)$ are both strictly positive (as indeed they must be if they are to form a John-Strömberg pair).

Throughout this proof we will let Q denote the d -dimensional unit cube, $Q = [0, 1]^d$. Let F_+ and F_- be two arbitrary disjoint subsets of Q which are both d -multi-cubes and satisfy

$$(3.1) \quad \min \{ \lambda(F_+), \lambda(F_-) \} > (1-\theta)\tau \lambda(Q \setminus F_+ \setminus F_-)$$

In view of Theorem 2.4, it will suffice to show that there exists a subcube W of $[0, 1]^d$ such that

$$(3.2) \quad \min \{ \lambda(W \cap F_+), \lambda(W \cap F_-) \} \geq (s - \theta(1-s)) \lambda(W).$$

For some sufficiently large integer N , both of the sets F_+ and F_- are finite unions of dyadic cubes all of the same side length 2^{-N} , as of course is the whole unit cube Q . Since the distance between F_+ and F_- must be positive, there must be some dyadic subcubes of Q of side length 2^{-N} which are not contained in $F_+ \cup F_-$, and thus their interiors are all contained in $Q \setminus F_+ \setminus F_-$. Therefore the set $Q \setminus F_+ \setminus F_-$ is also, at least to within some set of measure zero, a finite union of dyadic cubes of side length 2^{-N} .

Let $\mathcal{F}_+$ denote the collection of all dyadic cubes of side length 2^{-N} which are contained in F_+ . For each $\delta \in (0, 2^{-N-1})$ and each cube $W = \prod_{j=1}^n [a_j, a_j + 2^{-N}]$ in the collection $\mathcal{F}_+$, let $H(W, \delta)$ denote the cube concentric with W contained in the interior of W whose side length is $2^{-N} - 2\delta$, i.e., $H(W, \delta) = \prod_{j=1}^n [a_j + \delta, a_j + 2^{-N} - \delta]$. We will later

use the following obvious fact: If V is a cube which intersects with $H(W, \delta)$ and has side length less than δ , then

$$(3.3) \quad V \subset W.$$

We are going to choose a particular value of δ which will remain unchanged for the rest of this proof. Let $H_\delta^+ = \bigcup_{W \in \mathcal{F}_+} H(W, \delta)$. Obviously $\lambda(F_+) - \lambda(H_\delta^+) = \lambda(F_+ \setminus H_\delta^+)$ for each $\delta \in (0, 2^{-N-1})$ and $\lim_{\delta \searrow 0} \lambda(F_+ \setminus H_\delta^+) = 0$. Therefore, since the inequality in (3.1) is strict, we can and will choose our fixed value of δ to be sufficiently small to ensure that

$$(3.4) \quad \min \{ \lambda(H_\delta^+), \lambda(F_-) \} > (1 - \theta) \tau \lambda(Q \setminus F_+ \setminus F_-) + \tau \lambda(F_+ \setminus H_\delta^+).$$

Let $\mathcal{G}$ denote the collection of all dyadic cubes of side length 2^{-N} whose interiors are contained in $Q \setminus F_+ \setminus F_-$. Given an arbitrary positive integer k , we divide each cube in the collection $\mathcal{G}$ into 2^{dk} dyadic subcubes of side length 2^{-N-k} . Let $\mathcal{G}_k$ denote the collection of all dyadic cubes obtained in this way. In other words, $\mathcal{G}_k$ is simply the collection of all dyadic cubes of side length 2^{-N-k} whose interiors are contained in $Q \setminus F_+ \setminus F_-$. For each $W \in \mathcal{G}_k$ we let $U(W, k)$ be the (closed) cube concentric with W whose volume satisfies

$$(3.5) \quad \lambda(U(W, k)) = \theta \lambda(W).$$

Let $V_k = \bigcup_{W \in \mathcal{G}_k} U(W, k)$ and then let $H_k^- = F_- \cup V_k$.

In accordance with usual very standard notation, we will denote the interior and the boundary of any given cube W by W° and ∂W respectively. We of course have

$$(3.6) \quad Q \setminus F_+ \setminus F_- = \left(\bigcup_{W \in \mathcal{G}_k} W^\circ \right) \cup Z$$

for some set $Z \subset \bigcup_{W \in \mathcal{G}_k} \partial W$. Since $\lambda(Z) = 0$, this implies that

$$(3.7) \quad \lambda(Q \setminus F_+ \setminus F_-) = \sum_{W \in \mathcal{G}_k} \lambda(W).$$

In the following calculation we shall use (3.6) in the second line, and then, in the third line, the fact that, for every cube $W \in \mathcal{G}_k$, the set Z of zero measure introduced in (3.6) is disjoint from the cube W° and therefore also from $U(W, k)$. The final line of the calculation will use the facts that, for each pair of cubes W and W' in $\mathcal{G}_k$, we have $U(W, k) \subset W^\circ$ and also $W^\circ \cap (W')^\circ = \emptyset$ and therefore $W^\circ \cap U(W', k) = \emptyset$.

Thus we obtain that

$$\begin{aligned}
Q \setminus F_+ \setminus H_k^- &= Q \setminus F_+ \setminus F_- \setminus \left(\bigcup_{W \in \mathcal{G}_k} U(W, k) \right) \\
&= \left(Z \cup \left(\bigcup_{W \in \mathcal{G}_k} W^\circ \right) \right) \setminus \left(\bigcup_{W \in \mathcal{G}_k} U(W, k) \right) \\
&= Z \cup \left(\left(\bigcup_{W \in \mathcal{G}_k} W^\circ \right) \setminus \left(\bigcup_{W \in \mathcal{G}_k} U(W, k) \right) \right) \\
&= Z \cup \left(\bigcup_{W \in \mathcal{G}_k} (W^\circ \setminus U(W, k)) \right).
\end{aligned}$$

Since $\lambda(Z) = 0$, the preceding equalities, together with (3.5) and (3.7), imply that

$$\begin{aligned}
\lambda(Q \setminus F_+ \setminus H_k^-) &= \sum_{W \in \mathcal{G}_k} \lambda(W^\circ \setminus U(W, k)) = (1 - \theta) \sum_{W \in \mathcal{G}_k} \lambda(W) \\
(3.8) \qquad \qquad \qquad &= (1 - \theta) \lambda(Q \setminus F_+ \setminus F_-).
\end{aligned}$$

It is straightforward to check, using the facts that $F_+ \cap H_k^- = \emptyset$ and $H_\delta^+ \subset F_+$, that

$$(3.9) \qquad Q \setminus H_\delta^+ \setminus H_k^- = (Q \setminus F_+ \setminus H_k^-) \cup (F_+ \setminus H_\delta^+).$$

Since the two sets in parentheses on the right side of (3.9) are disjoint, it follows that

$$(3.10) \qquad \lambda(Q \setminus F_+ \setminus H_k^-) + \lambda(F_+ \setminus H_\delta^+) = \lambda(Q \setminus H_\delta^+ \setminus H_k^-).$$

Now we can first use the fact that $F_- \subset H_k^-$ and then invoke (3.4) followed by (3.8) and then (3.10), to obtain that

$$\begin{aligned}
\min \{ \lambda(H_\delta^+), \lambda(H_k^-) \} &\geq \min \{ \lambda(H_\delta^+), \lambda(F_-) \} \\
&> (1 - \theta) \tau \lambda(Q \setminus F_+ \setminus F_-) + \tau \lambda(F_+ \setminus H_\delta^+) \\
&= \tau \lambda(Q \setminus F_+ \setminus H_k^-) + \tau \lambda(F_+ \setminus H_\delta^+) \\
(3.11) \qquad \qquad \qquad &= \tau \lambda(Q \setminus H_\delta^+ \setminus H_k^-).
\end{aligned}$$

Since $V_k \subset Q \setminus F_+ \setminus F_-$ and $F_+ \cap F_- = \emptyset$, we have

$$(3.12) \qquad H_k^- \subset F_- \cup (Q \setminus F_+ \setminus F_-) = Q \setminus F_+.$$

Furthermore $H_\delta^+ \subset F_+$ and so H_δ^+ and H_k^- are disjoint measurable subsets of Q . According to our hypotheses, (τ, s) is a John-Strömberg pair for $Q(\mathbb{R}^d)$. Therefore, by the estimates leading to (3.11), for each $k \in \mathbb{N}$, there must exist some subcube W_k of Q such that

$$(3.13) \qquad \min \{ \lambda(W_k \cap H_\delta^+), \lambda(W_k \cap H_k^-) \} \geq s \lambda(W_k).$$

We now claim that the side length of W_k , which we can conveniently write as $(\lambda(W_k))^{1/d}$, must satisfy

$$(3.14) \qquad (\lambda(W_k))^{1/d} \geq \delta.$$

To show this we first observe that, since the cube W_k intersects with H_δ^+ , it must intersect with the cube $H(W, \delta)$ for at least one cube W in the collection $\mathcal{F}_+$. If (3.14) does not hold, i.e., if W_k has side length less than δ , then (cf. the discussion immediately preceding (3.3)) W_k must be completely contained in that particular cube W and therefore also in F_+ . Consequently W_k cannot intersect with the set H_k^- . (Here we have used (3.12) once more.) This contradicts (3.13) and shows that (3.14) does hold.

Let $\widetilde{W}_k$ be a new cube containing W_k and concentric with W_k . More precisely, if $W_k = \prod_{j=1}^d [a_j, b_j]$, then we choose $\widetilde{W}_k$ to be $\prod_{j=1}^d [a_j - 2^{-N-k}, b_j + 2^{-N-k}]$. Clearly $\widetilde{W}_k$ contains all dyadic cubes of side length 2^{-N-k} which intersect with W_k , and the side length $\left(\lambda(\widetilde{W}_k)\right)^{1/d}$ of $\widetilde{W}_k$ satisfies

$$\left(\lambda(\widetilde{W}_k)\right)^{1/d} = (\lambda(W_k))^{1/d} + 2^{1-N-k}.$$

This, together with (3.14), gives us that

$$\begin{aligned} \left(\frac{\lambda(\widetilde{W}_k)}{\lambda(W_k)}\right)^{1/d} &= \frac{(\lambda(W_k))^{1/d} + 2^{1-N-k}}{(\lambda(W_k))^{1/d}} \\ &= 1 + \frac{2^{1-N-k}}{(\lambda(W_k))^{1/d}} \leq 1 + \frac{2^{1-N-k}}{\delta}. \end{aligned}$$

It follows that

$$\begin{aligned} \lambda(\widetilde{W}_k) - \lambda(W_k) &= \lambda(W_k) \left(\frac{\lambda(\widetilde{W}_k)}{\lambda(W_k)} - 1 \right) \\ (3.15) \qquad \qquad &\leq \lambda(W_k) \left(\left(1 + \frac{2^{1-N-k}}{\delta} \right)^d - 1 \right). \end{aligned}$$

Let $\mathcal{H}_k$ be the collection of all cubes in $\mathcal{G}_k$ which intersect with W_k . Clearly

$$(3.16) \qquad \bigcup_{W \in \mathcal{H}_k} W^\circ \subset (Q \setminus F_+ \setminus F_-) \cap \widetilde{W}_k.$$

Our definitions of V_k and of $\mathcal{H}_k$ also immediately give us that

$$\begin{aligned} W_k \cap V_k &= W_k \cap \left(\bigcup_{W \in \mathcal{G}_k} U(W, k) \right) \\ &= W_k \cap \left(\bigcup_{W \in \mathcal{H}_k} U(W, k) \right) \\ &\subset \bigcup_{W \in \mathcal{H}_k} U(W, k). \end{aligned}$$

This will imply the first line in the following calculation. Its second line will use the fact that the interiors of the cubes in $\mathcal{G}_k$ and therefore also in $\mathcal{H}_k$, are pairwise disjoint. Its third line will use (3.16). Its fourth line will use the fact that $H_\delta^+ \subset F_+$. Its sixth line will follow from the obvious inclusion $W_k \subset \widetilde{W}_k$. The justifications of all other steps should be evident. (We wonder, casually, whether it might somehow be possible to replace the simple-minded transition from the sixth to the seventh line by a sharper estimate, which might then lead to a (slightly) stronger version of Theorem 3.1.)

$$\begin{aligned}
\lambda(W_k \cap V_k) &\leq \sum_{W \in \mathcal{H}_k} \lambda(U(W, k)) = \theta \sum_{W \in \mathcal{H}_k} \lambda(W) \\
&= \theta \sum_{W \in \mathcal{H}_k} \lambda(W^\circ) = \theta \lambda \left(\bigcup_{W \in \mathcal{H}_k} W^\circ \right) \\
&\leq \theta \lambda \left((Q \setminus F_+ \setminus F_-) \cap \widetilde{W}_k \right) \\
&\leq \theta \lambda \left((Q \setminus H_\delta^+ \setminus F_-) \cap \widetilde{W}_k \right) \\
&\leq \theta \lambda \left((Q \setminus H_\delta^+ \setminus F_-) \cap W_k \right) + \theta \lambda \left(\widetilde{W}_k \setminus W_k \right) \\
&= \theta \lambda \left((Q \setminus H_\delta^+ \setminus F_-) \cap W_k \right) + \theta \left(\lambda \left(\widetilde{W}_k \right) - \lambda(W_k) \right) \\
&\leq \theta \lambda \left((Q \setminus H_\delta^+) \cap W_k \right) + \theta \left(\lambda \left(\widetilde{W}_k \right) - \lambda(W_k) \right) \\
&= \theta \lambda \left(W_k \setminus H_\delta^+ \right) + \theta \left(\lambda \left(\widetilde{W}_k \right) - \lambda(W_k) \right).
\end{aligned}$$

Combining the result of this calculation with (3.15), we deduce that, for each $k \in \mathbb{N}$,

$$(3.17) \quad \lambda(W_k \cap V_k) \leq \theta \lambda \left(W_k \setminus H_\delta^+ \right) + \varepsilon_k \lambda(W_k)$$

where the sequence of positive numbers $\{\varepsilon_k\}_{k \in \mathbb{N}} := \left\{ \theta \left(\left(1 + \frac{2^{1-N-k}}{\delta} \right)^d - 1 \right) \right\}_{k \in \mathbb{N}}$ satisfies

$$(3.18) \quad \lim_{k \rightarrow \infty} \varepsilon_k = 0.$$

In view of (3.13), we have that

$$(3.19) \quad \lambda(W_k \setminus H_\delta^+) = \lambda(W_k) - \lambda(W_k \cap H_\delta^+) \leq (1 - s) \lambda(W_k).$$

We now have the ingredients needed to estimate $\lambda(W_k \cap F_-)$ from below. Since $\lambda(W_k \cap H_k^-) = \lambda(W_k \cap F_-) + \lambda(W_k \cap V_k)$ we can use (3.13) and (3.17) and then (3.19) to obtain that

$$\begin{aligned}
\lambda(W_k \cap F_-) &= \lambda(W_k \cap H_k^-) - \lambda(W_k \cap V_k) \\
&\geq s \lambda(W_k) - \theta \lambda \left(W_k \setminus H_\delta^+ \right) - \varepsilon_k \lambda(W_k) \\
&= s \lambda(W_k) - \theta (1 - s) \lambda(W_k) - \varepsilon_k \lambda(W_k) \\
(3.20) \quad &= (s - \theta(1 - s) - \varepsilon_k) \lambda(W_k).
\end{aligned}$$

Furthermore, again with the help of (3.13), we also obviously have that

$$(3.21) \quad \lambda(W_k \cap F_+) \geq \lambda(W_k \cap H_\delta^+) \geq s\lambda(W_k).$$

As already observed at the beginning of this proof, $s - \theta(1 - s)$ and $\theta(1 - s)$ are both strictly positive. So $s > (s - \theta(1 - s) - \varepsilon_k)$ and $(s - \theta(1 - s) - \varepsilon_k) > 0$ for all sufficiently large k . Therefore, we can now deduce from (3.20) and (3.21) that $((1 - \theta)\tau, s - \theta(1 - s) - \varepsilon_k)$ is a John-Strömberg pair for $\mathcal{Q}(\mathbb{R}^d)$ for all sufficiently large k . This seems to be very close to our required result. Indeed it will only require a little more effort to obtain that result.

For each $k \in \mathbb{N}$, let x_k be the centre of the cube W_k and let r_k be half its side length. I.e., we can set $W_k = Q(x_k, r_k)$ in the standard notation recalled above in Definition 1.1(ii). Of course $x_k \in Q$ and, by (3.14) and the fact that $W_k \subset Q$, we also have $\delta/2 \leq r_k \leq 1/2$. Therefore there exists a strictly increasing sequence $\{n_k\}_{k \in \mathbb{N}}$ of positive integers such that the sequences $\{x_{n_k}\}_{k \in \mathbb{N}}$ and $\{r_{n_k}\}_{k \in \mathbb{N}}$ converge, respectively, to a point $x \in Q$ and a number $r \in [\delta/2, 1/2]$. Let W be the cube $W = Q(x, r)$. Then, by Lemma 10.1 of [1, p. 43] and (3.20) and (3.18), it follows that $W \subset Q$ and

$$\begin{aligned} \lambda(W \cap F_-) &= \lim_{k \rightarrow \infty} \lambda(W_{n_k} \cap F_-) \\ &\geq \lim_{k \rightarrow \infty} (s - \theta(1 - s) - \varepsilon_{n_k}) \lambda(W_{n_k}) \\ &= (s - \theta(1 - s)) \lambda(W). \end{aligned}$$

Similarly, replacing k by n_k in (3.21) and passing to the limit as k tends to ∞ , we obtain that

$$\lambda(W \cap F_+) \geq s\lambda(W) \geq (s - \theta(1 - s))\lambda(W).$$

All this shows that the subcube W of Q satisfies (3.2) and therefore completes the proof of the theorem. $\square$

Remark 3.3. It seems quite possible that a more elaborate version of the preceding proof might show that the hypotheses of Theorem 3.1 can yield a stronger result, namely that (τ', s') is a John-Strömberg pair for some positive number τ' smaller than $(1 - \theta)\tau$ and/or for some number s' greater than $s - \theta(1 - s)$. Perhaps a strategy for proving this might involve separately considering the two cases where $\lambda(F_+) > c\lambda(F_-)$ and where $\lambda(F_-) \leq \lambda(F_+) \leq c\lambda(F_-)$ for some suitably chosen constant $c > 1$.

4. SOME FURTHER RESULTS ABOUT JOHN-STRÖMBERG PAIRS

4.1. Other characterizations of John-Strömberg pairs. It will be convenient to begin by introducing some “technical” terminology.

Definition 4.1. Let F_+ and F_- be two disjoint measurable subsets of $[0, 1]^d$ which both have positive measure and which also satisfy one of the following two conditions:

- (i) $\lambda([0, 1]^d \setminus F_+ \setminus F_-) = 0$.
- (ii) There exists a non-empty open subset Ω of $[0, 1]^d$ for which

$$\max \{ \lambda(F_+ \cap \Omega), \lambda(F_- \cap \Omega) \} < \lambda(\Omega) \quad \text{and} \quad \min \{ \lambda(F_+ \cap \Omega), \lambda(F_- \cap \Omega) \} = 0.$$

Then we shall say that (F_+, F_-) is a *tame couple* in $[0, 1]^d$.

Remark 4.2. Obviously the two conditions (i) and (ii) are mutually exclusive.

We can now present the new variant of Theorem 2.4 to which we referred in the preamble to that theorem. It specifies two more properties of a pair (τ, s) , which we label as $(**)$ and $(***)$, and which are each equivalent to the property that $(\tau, s) \in JS(d)$. Note that the only difference between the statement of property $(**)$ in Theorem 4.3, and the statement of property $(*)$ in Theorem 2.4 is that the strict equality “ $>$ ” which appears in (2.3) in the statement of $(*)$ has been replaced by “ $\geq$ ” in (4.1) in the statement of $(**)$. Property $(***)$ of Theorem 4.3 is more elaborate, and requires the terminology of Definition 4.1.

Theorem 4.3. *Let d be a positive integer and let (τ, s) be an ordered pair of positive numbers. Then each of the following two properties is equivalent to the property that (τ, s) is a John-Strömberg pair for $\mathcal{Q}(\mathbb{R}^d)$.*

*(**) Whenever F_+ and F_- are disjoint d -multi-cubes which satisfy*

$$(4.1) \quad \min \{ \lambda(F_+), \lambda(F_-) \} \geq \tau \lambda([0, 1]^d \setminus F_+ \setminus F_-),$$

then there exists a cube W contained in $[0, 1]^d$ for which

$$(4.2) \quad \min \{ \lambda(W \cap F_+), \lambda(W \cap F_-) \} \geq s \lambda(W).$$

*(***) Whenever (F_+, F_-) is a tame couple in $[0, 1]^d$ which satisfies (4.1), then there exists a cube W contained in $[0, 1]^d$ for which (4.2) holds.*

Proof. Let us first show that property $(***)$ implies property $(**)$. Suppose that F_+ and F_- are arbitrary disjoint d -multi-cubes. By the same reasoning as was given in the paragraph immediately following (3.2) in the proof of Theorem 3.1, we see that the set $[0, 1]^d \setminus F_+ \setminus F_-$ must contain the interior Ω of at least one dyadic subcube of $[0, 1]^d$. Such a set Ω of course satisfies the condition (ii) in Definition 4.1. Consequently, (F_+, F_-) is a tame couple in $[0, 1]^d$. It immediately follows that property $(***)$ indeed does imply property $(**)$.

Obviously, if the pair (τ, s) of positive numbers has property $(**)$, then it also has property $(*)$ of Theorem 2.4 and therefore (τ, s) is a John-Strömberg pair for $\mathcal{Q}(\mathbb{R}^d)$.

Now suppose that (τ, s) is a John-Strömberg pair for $\mathcal{Q}(\mathbb{R}^d)$. Then, by letting $\{\tau_n\}_{n \in \mathbb{N}}$ and $\{s_n\}_{n \in \mathbb{N}}$ be the special constant sequences $\tau_n = \tau$ and $s_n = s$, we see that (τ, s) satisfies the hypotheses of Lemma 4.5 which we will formulate and prove below. In view of part (b) of that lemma, (τ, s) has property $(***)$. This completes the proof of the theorem. $\square$

Remark 4.4. It is very natural to wonder whether property $(***)$ is equivalent to a stronger and more simply expressed variant of that property in which the same implication is required to hold for *all* pairs of disjoint measurable subsets F_+, F_- of $[0, 1]^d$ which both have positive measure, thus omitting the requirement that F_+ and F_- should form a tame couple. (I.e., as in Remark 7.17 of [1, p. 30], we are essentially wondering whether in Definition 2.1, at least in the case where $\mathcal{E} = \mathcal{Q}(\mathbb{R}^d)$, it would be equivalent to replace “ $>$ ” by “ $\geq$ ” in (2.1), of course then with the necessary proviso that $\lambda(E_+)$ and $\lambda(E_-)$ are both positive.) We are unable to answer this question, but Lemma 4.5 shows that if its answer is negative, then the sets F_+ and F_- which provide a counterexample,

must both have very intricate structure (and it would not be inappropriate to refer to them as forming a “wild” couple).

Lemma 4.5. *Let d be a positive integer. Let $\{\tau_n\}_{n \in \mathbb{N}}$ and $\{s_n\}_{n \in \mathbb{N}}$ be two sequences of positive numbers which converge to positive limits, τ and s respectively. Suppose that (τ_n, s_n) is a John-Strömberg pair for $\mathcal{Q}(\mathbb{R}^d)$ for every $n \in \mathbb{N}$. Then*

- (a) *the limiting pair (τ, s) has the property $(**)$ of Theorem 4.3, and,*
- (b) *in the special case where $s_n \geq s$ for every $n \in \mathbb{N}$, the limiting pair (τ, s) also has the property $(***)$ of Theorem 4.3.*

Proof. Let us fix an arbitrary $d \in \mathbb{N}$, and arbitrary sequences $\{\tau_n\}_{n \in \mathbb{N}}$ and $\{s_n\}_{n \in \mathbb{N}}$ of positive numbers which tend respectively to the positive numbers τ and s , and have the property that $(\tau_n, s_n) \in JS(d)$ for each $n \in \mathbb{N}$.

In order to deduce that (τ, s) has property $(**)$ and/or $(***)$, we begin by fixing two arbitrary disjoint measurable sets F_+ and F_- which are contained in $[0, 1]^d$, which form a tame couple in $[0, 1]^d$, and which also satisfy (4.1). To obtain part (a) of the lemma we have the task of proving, for these choices of F_+ , F_- , $\{\tau_n\}_{n \in \mathbb{N}}$, $\{s_n\}_{n \in \mathbb{N}}$, τ and s , that there exists a subcube W of $[0, 1]^d$ which satisfies (4.2). If needed at any stage of our proof of that fact, we may make the additional assumption that F_+ and F_- are both d -multi-cubes. An analogous task is required to obtain part (b) of the lemma, with the difference that in our proof this time, instead of being able to assume that F_+ and F_- are d -multi-cubes, we may make the additional assumption, if needed, that $s_n \geq s$ for all n .

There is quite a lot of overlap in the ingredients which will be used for performing these two tasks, and it may help avoid some confusion if we give some general descriptions, in advance, of the four steps which we shall use to accomplish both of them, almost simultaneously. The two above mentioned additional assumptions will not be needed in the first two of these steps.

In Step 1 of the proof, we shall use the sets F_+ and F_- and the sequences $\{\tau_n\}_{n \in \mathbb{N}}$ and $\{s_n\}_{n \in \mathbb{N}}$ to construct a special sequence $\{W(n_k)\}_{k \in \mathbb{N}}$ of subcubes of $[0, 1]^d$ which has several convenient properties. In particular, this sequence converges in an obvious way (which we will explicitly define below) either to a limiting cube or to a point.

In Step 2, we shall see that if the sequence $\{W(n_k)\}_{k \in \mathbb{N}}$ obtained in the preceding step converges to a subcube W , then that cube is contained in $[0, 1]^d$ and satisfies (4.2).

In Step 3, we shall see that whenever the disjoint measurable sets F_+ and F_- are both required to also be d -multi-cubes, then the sequence $\{W(n_k)\}_{k \in \mathbb{N}}$ can be always be constructed so that it converges to a subcube. In view of Step 2, this will complete the proof of part (a) of the lemma.

Finally, in Step 4, in order to complete the proof of part (b), it will remain (again in view of Step 2) only to deal with the case where the sequence $\{W(n_k)\}_{k \in \mathbb{N}}$ converges to a point. We will do this by showing that, in this case, when we also impose the requirement that the sequence $\{s_n\}_{n \in \mathbb{N}}$ satisfies $s_n \geq s$ for each n , then there exists a positive integer k_0 for which the subcube $W = W(n_{k_0})$ is contained in $[0, 1]^d$ and satisfies (4.2).

STEP 1: Construction and properties of the special sequence $\{W(n_k)\}_{k \in \mathbb{N}}$.

Since (F_+, F_-) is a tame couple, it must satisfy either condition (i) or condition (ii) of Definition 4.1. The construction of $\{W(n_k)\}_{k \in \mathbb{N}}$ is very simple when condition (i) holds, but we shall defer its description to later, and first consider the case when F_+ and F_- satisfy condition (ii).

In that case, since the roles of F_+ and F_- are interchangeable in properties (**) and (***) and in Definition 4.1 and in (4.2), we can assume, without loss of generality, that there exists some open subset Ω of $[0, 1]^d$ for which

$$(4.3) \quad 0 = \lambda(F_- \cap \Omega) \leq \lambda(F_+ \cap \Omega) < \lambda(\Omega).$$

Of course Ω must be non-empty. By Theorem 1.11 of [10, p. 8], Ω can be expressed as the union of a sequence of non-overlapping dyadic cubes. It follows from (4.3) that at least one of these cubes, which we will denote by V , must satisfy $0 \leq \lambda(F_+ \cap V) < \lambda(V)$ and of course also $\lambda(F_- \cap V) = 0$. Since

$$\lambda(V^\circ \setminus F_+) = \lambda(V \setminus F_+) = \lambda(V) - \lambda(V \cap F_+) > 0,$$

(where, as usual, V° denotes the interior of V), the Lebesgue differentiation theorem guarantees the existence of a point $z \in V^\circ \setminus F_+$ for which $\lim_{r \searrow 0} \frac{\lambda(Q(z, r) \cap V^\circ \setminus F_+)}{\lambda(Q(z, r))} = 1$ and, consequently, $\lim_{r \searrow 0} \frac{\lambda(Q(z, r) \cap F_+)}{\lambda(Q(z, r))} = 0$. In particular, this enables us to assert the existence of a sequence $\{\rho_k\}_{k \in \mathbb{N}}$ of positive numbers tending to monotonically to zero, such that $Q(z, \rho_k) \subset V^\circ$ and $\lambda(Q(z, \rho_k) \cap F_+) < \lambda(Q(z, \rho_k))$ for each $k \in \mathbb{N}$.

Let us now define a sequence $\{F_+(k)\}_{k \in \mathbb{N}}$ of measurable sets by setting

$$(4.4) \quad F_+(k) = F_+ \cup Q(z, \rho_k) \text{ for each } k \in \mathbb{N}.$$

We also introduce the set $G := F_- \setminus V^\circ$. Note that $F_+(k)$ and G are disjoint for each $k \in \mathbb{N}$. In fact

$$\begin{aligned} F_+(k) \cap G &= (F_+ \cup Q(z, \rho_k)) \cap G = (F_+ \cap G) \cup (Q(z, \rho_k) \cap G) \\ &\subset (F_+ \cap F_-) \cup (Q(z, \rho_k) \setminus V^\circ) = \emptyset \cup \emptyset. \end{aligned}$$

The properties of the sequence $\{\rho_k\}_{k \in \mathbb{N}}$ ensure that $\lambda(F_+(k)) > \lambda(F_+)$ for each k . Obviously $V^\circ \setminus F_+(k) = V^\circ \setminus F_+(k) \setminus G$. Consequently,

$$\begin{aligned} &\lambda([0, 1]^d \setminus F_+(k) \setminus G) \geq \lambda(V^\circ \setminus F_+(k) \setminus G) \\ &= \lambda(V^\circ \setminus F_+(k)) \geq \lambda(Q(z, \rho_k) \setminus F_+(k)) = \lambda(Q(z, \rho_k)) - \lambda(Q(z, \rho_k) \cap F_+(k)) > 0. \end{aligned}$$

We also have

$$(4.5) \quad \lambda(F_-) = \lambda(F_- \cap V^\circ) + \lambda(F_- \setminus V^\circ) = \lambda(F_- \setminus V^\circ) = \lambda(G).$$

Using the various properties of $F_+(k)$ and G established above, we see that

$$\begin{aligned} 0 &< \lambda([0, 1]^d \setminus F_+(k) \setminus G) = 1 - \lambda(F_+(k)) - \lambda(G) \\ &< 1 - \lambda(F_+) - \lambda(F_-) = \lambda([0, 1]^d \setminus F_+ \setminus F_-). \end{aligned}$$

Since $\tau > 0$, this implies that

$$(4.6) \quad 0 < \tau \lambda([0, 1]^d \setminus F_+(k) \setminus G) < \tau \lambda([0, 1]^d \setminus F_+ \setminus F_-).$$

Since $\min \{\lambda(F_+(k)), \lambda(G)\} \geq \min \{\lambda(F_+), \lambda(F_-)\}$, it follows from (4.6) and the fact that F_+ and F_- satisfy (4.1) that

$$\min \{\lambda(F_+(k)), \lambda(G)\} > \tau \lambda([0, 1]^d \setminus F_+(k) \setminus G) > 0 \text{ for each } k \in \mathbb{N}.$$

We use these strict inequalities to construct a sequence $\{j(k)\}_{k \in \mathbb{N}}$ of positive integers such that $j(k) \geq k$ and $j(k)$ is sufficiently large to ensure that $\tau_{j(k)}$ is sufficiently close to τ to imply that

$$\min \{\lambda(F_+(k)), \lambda(G)\} > \tau_{j(k)} \lambda([0, 1]^d \setminus F_+(k) \setminus G) \text{ for each } k \in \mathbb{N}.$$

Since $(\tau_{j(k)}, s_{j(k)}) \in JS(d)$ and since $F_+(k)$ and G are disjoint, we see, in accordance with Definition 2.1 (for $\mathcal{E} = \mathcal{Q}(\mathbb{R}^d)$), that there exists a subcube $W(k)$ of $[0, 1]^d$ which satisfies

$$(4.7) \quad \min \{\lambda(W(k) \cap F_+(k)), \lambda(W(k) \cap G)\} \geq s_{j(k)} \lambda(W(k)).$$

The fact that $j(k) \geq k$ for each k ensures that

$$(4.8) \quad \lim_{k \rightarrow \infty} s_{j(k)} = s.$$

Since

$$W(k) \cap F_+ \subset W(k) \cap F_+(k) = (W(k) \cap F_+) \cup (W(k) \cap Q(z, r_k)) \subset (W(k) \cap F_+) \cup Q(z, r_k)$$

and $\lambda(Q(z, r_k)) = (2r_k)^d$, we obtain that

$$(4.9) \quad \lambda(W(k) \cap F_+) \leq \lambda(W(k) \cap F_+(k)) \leq \lambda(W(k) \cap F_+) + (2\rho_k)^d \text{ for all } k \in \mathbb{N}.$$

A slight variant of the simple reasoning in (4.5) gives us that

$$(4.10) \quad \lambda(W(k) \cap F_-) = \lambda(W(k) \cap G) \text{ for all } k \in \mathbb{N}.$$

We use the standard notation of Definition 1.1 (ii) and write $W(k) = Q(x_k, r_k)$, i.e., we let $x_k \in [0, 1]^d$ and $r_k > 0$ be the centre and half-side length respectively of the cube $W(k)$. There exists a strictly increasing sequence $\{n_k\}_{k \in \mathbb{N}}$ of positive integers such that the sequences $\{x_{n_k}\}_{k \in \mathbb{N}}$ and $\{r_{n_k}\}_{k \in \mathbb{N}}$ converge, respectively, to a point $x \in [0, 1]^d$ and a number $r \in [0, 1/2]$. This gives us the sequence $\{W(n_k)\}_{k \in \mathbb{N}} = \{Q(x_{n_k}, r_{n_k})\}_{k \in \mathbb{N}}$ which we have sought to construct in this step of the proof. We can now explicitly give the obvious definition of what we mean by the convergence of this sequence, namely that it converges to the cube $Q(x, r)$ if $r > 0$, and it converges to the point x if $r = 0$.

It still remains to construct this sequence in the (much simpler) case where F_+ and F_- satisfy condition (i) of Definition 4.1, i.e., when $\lambda([0, 1]^d \setminus F_+ \setminus F_-) = 0$. Here we can use Lemma 7.1 of [1, p. 25] or of [3, p. 150] to ensure the existence of a cube $W = Q(x, r) \subset [0, 1]^d$ for which

$$(4.11) \quad \lambda(W \cap F_+) = \lambda(W \setminus F_+) = \frac{1}{2} \lambda(W).$$

We also note that

$$(4.12) \quad \lambda(W \cap F_-) = \lambda(W \setminus F_+).$$

In this case we simply define $x_k = x$, $r_k = r$, $W(k) = W$ and $n_k = k$ for each $k \in \mathbb{N}$, and so the constant sequence $\{W(n_k)\}_{k \in \mathbb{N}}$ of course converges to the cube which is its constant value.

STEP 2: A proof that whenever the limit of $\{W(n_k)\}_{k \in \mathbb{N}}$ is a cube, then that cube has the two properties required to immediately complete the proof of the theorem.

Suppose that the limit of the sequence $\{W(n_k)\}_{k \in \mathbb{N}} = \{Q(x_{n_k}, r_{n_k})\}_{k \in \mathbb{N}}$ introduced in the previous step, is indeed a cube $W = Q(x, r)$, i.e., that

$$(4.13) \quad r := \lim_{k \rightarrow \infty} r_{n_k} > 0.$$

Let us now prove that this implies that W is contained in $[0, 1]^d$ and that it satisfies (4.2).

In the case which was dealt with at the very end of the previous step, where F_+ and F_- satisfy condition (i) of Definition 4.1, and where indeed we always have $r > 0$, it is already known that the cube W is contained in $[0, 1]^d$. We note that, by the reasoning in Remark 2.2, we have $s_n \leq 1/2$ for each $n \in \mathbb{N}$ and therefore also $s \leq 1/2$. So (4.2) follows immediately from (4.11) and (4.12).

We turn to the remaining case, where F_+ and F_- satisfy condition (ii) of Definition 4.1, and therefore the sequence $\{W(n_k)\}_{k \in \mathbb{N}}$ is constructed in the more elaborate way described in the first part of the previous step. Here the positivity of the limit r permits us to apply Lemma 10.1 of [1, p. 43] to the sequence $\{W(n_k)\}_{k \in \mathbb{N}} = \{Q(x_{n_k}, r_{n_k})\}_{k \in \mathbb{N}}$ to obtain the first required conclusion, that $W \subset [0, 1]^d$, and to also obtain that

$$(4.14) \quad \lambda(W \cap F_+) = \lim_{k \rightarrow \infty} \lambda(W(n_k) \cap F_+)$$

and, (here also using (4.10)), that

$$(4.15) \quad \lambda(W \cap F_-) = \lambda(W \cap G) = \lim_{k \rightarrow \infty} \lambda(W(n_k) \cap G),$$

and also that

$$(4.16) \quad \lambda(W) = \lim_{k \rightarrow \infty} \lambda(W(n_k)).$$

From (4.14) and (4.9) we see that

$$(4.17) \quad \lambda(W \cap F_+) = \lim_{k \rightarrow \infty} \lambda(W(n_k) \cap F_+(n_k)),$$

and from (4.16) and (4.8) we see that

$$(4.18) \quad s\lambda(W) = \lim_{k \rightarrow \infty} s_{j(n_k)}\lambda(W(n_k)).$$

In view of (4.7) we have that

$$\lambda(W(n_k) \cap F_+(n_k)) \geq s_{j(n_k)}\lambda(W(n_k)) \quad \text{and} \quad \lambda(W(n_k) \cap G) \geq s_{j(n_k)}\lambda(W(n_k))$$

for each $k \in \mathbb{N}$. If we take the limit as k tends to ∞ in each of these two inequalities, and apply (4.17), (4.15) and (4.18), then the two resulting inequalities can be rewritten as the required single inequality (4.2).

Thus we have shown that in both cases, i.e., whether it is condition (i) or condition (ii) of Definition 4.1 which applies to F_+ and F_- , if (4.13) holds, then the limiting cube W indeed has both the properties required to complete the proof of part (a) and also part (b) of the lemma. It remains to explain how we can sometimes guarantee that (4.13) does hold, and how we can proceed when it does not hold.

STEP 3: Completion of the proof of part (a) of the lemma.

Our preceding treatment of the case where (4.13) holds, will now enable us to complete the proof of part (a) of the lemma in full generality. Part (a) refers to the property (**), so in our proof of it we can and must assume that F_+ and F_- are both d -multi-cubes. Once again we refer to the reasoning at the beginning of Theorem 4.3 which shows that F_+ and F_- necessarily satisfy condition (ii) of Definition 4.1 and enables us to choose an open set Ω which satisfies (4.3) and is the interior of a dyadic interval contained in $[0, 1]^d \setminus F_+ \setminus F_-$. Then in fact we can also simply take $V^\circ = \Omega$. This choice gives us that the set G introduced in preceding steps of our proof, must satisfy $G = F_- \setminus V^\circ = F_-$.

Since they are d -multi-cubes, F_+ and $F_- = G$ are compact and so of course is $Q(z, \rho_k)$. Consequently, the set $F_+(k)$ (defined by (4.4)) is also compact. The distance, which we denote by $\text{dist}_\infty(F_+(k), G)$, between the disjoint compact sets $F_+(k)$ and G with respect to the ℓ^∞ metric on $\mathbb{R}^d$, must be positive. In fact, since $F_+(k) \subset F_+(1)$ (because $\rho_k \leq \rho_1$), we have that

$$(4.19) \quad \text{dist}_\infty(F_+(k), G) \geq \text{dist}_\infty(F_+(1), G) > 0 \text{ for all } k \in \mathbb{N}.$$

Now we proceed more or less similarly to the last steps of the proof of Theorem 3.1. Since the cube $W(k) = Q(x_k, r_k)$ intersects both of the sets $F_+(k)$ and G , its side length $2r_k$ cannot be smaller than $\text{dist}_\infty(F_+(k), G)$. Consequently, also using (4.19), we see that

$$\frac{1}{2} \geq r_k \geq \delta_0 := \frac{1}{2} \text{dist}_\infty(F_+(1), G) > 0 \text{ for all } k \in \mathbb{N}.$$

Consequently $r \geq \delta_0$ and so (4.13) holds, permitting us to use the reasoning of Step 2 to ensure the existence of cube W which has the properties required to complete the proof of part (a).

STEP 4: Completion of the proof of part (b) of the lemma.

In view of Step 2, if the cubes $W(n_k)$ converge to the cube W , then that cube is contained in $[0, 1]^d$ and satisfies (4.2) and no further reasoning is required to complete the proof of part (b). Thus it remains only to deal with the case where $r = \lim_{k \rightarrow \infty} r_{n_k} = 0$. Let us first see that in this case the point $x = \lim_{k \rightarrow \infty} x_{n_k}$ cannot coincide with the point z which appears in the definition (4.4) of the sets $F_+(k)$. If $x = z$ and is therefore in V° , then there exists some k which is sufficiently large to ensure that $r_{n_k} + \|x_{n_k} - x\|_\infty < \text{dist}_\infty(x, \partial V)$. This means that every point y in the cube $W(n_k) = Q(x_{n_k}, r_{n_k})$ satisfies $\|y - x\|_\infty \leq \|y - x_{n_k}\|_\infty + \|x_{n_k} - x\|_\infty < \text{dist}_\infty(x, \partial V)$. Consequently $W(n_k) \subset V^\circ$ and therefore $W(n_k) \cap G = \emptyset$. Since this contradicts (4.7), we have indeed shown that $x \neq z$.

In view of this fact, there exists an integer k_0 which is large enough to ensure that

$$(4.20) \quad \|x - x_{n_{k_0}}\| + \rho_{n_{k_0}} + r_{n_{k_0}} < \|x - z\|_\infty,$$

where $\rho_{n_{k_0}}$ is a element of the sequence $\{\rho_k\}_{k \in \mathbb{N}}$ with limit 0 which is used in the definition (4.4) of the sets $F_+(k)$. If y is a point in the intersection of the two cubes

$W(n_{k_0}) = Q(x_{n_{k_0}}, r_{n_{k_0}})$ and $Q(z, \rho_{n_{k_0}})$, then

$$\begin{aligned} \|x - z\|_\infty &\leq \|x - x_{n_{k_0}}\| + \|x_{n_{k_0}} - z\|_\infty \leq \|x - x_{n_{k_0}}\| + \|x_{n_{k_0}} - y\|_\infty + \|y - z\|_\infty \\ &\leq \|x - x_{n_{k_0}}\| + \rho_{n_{k_0}} + r_{n_{k_0}}. \end{aligned}$$

But this contradicts (4.20) and enables us to conclude that $W(n_{k_0})$ and $Q(z, \rho_{n_{k_0}})$ must be disjoint and therefore that

$$W(n_{k_0}) \cap F_+(n_{k_0}) = W(n_{k_0}) \cap F_+.$$

We apply this, together with (4.10) for $k = n_{k_0}$ and then (4.7) for $k = n_{k_0}$, to obtain that the cube $W := W(n_{k_0})$ satisfies

$$\begin{aligned} \min \{ \lambda(W \cap F_+), \lambda(W \cap F_-) \} &= \min \{ \lambda(W \cap F_+(n_{k_0})), \lambda(W \cap G_-) \} \\ (4.21) \qquad \qquad \qquad &\geq s_{j(n_{k_0})} \lambda(W). \end{aligned}$$

The cube W , like all other cubes in the sequence $\{W(n_k)\}_{k \in \mathbb{N}}$ is contained in $[0, 1]^d$. Finally we have to recall that in the statement of part (b) of the lemma, the sequence $\{s_n\}_{n \in \mathbb{N}}$ is required to satisfy $s_n \geq s$ and so (4.21) shows that, also in this last remaining case, we have obtained a subcube W of $[0, 1]^d$ which satisfies (4.2) for the given sets F_+ and F_- . This completes the proof of the lemma. $\square$

4.2. Some pairs which are not John-Strömberg pairs. The following result is a more elaborate variant of Remark 2.2.

Lemma 4.6. *For each $d \in \mathbb{N}$ and each $\tau > 0$, and for every $s > 1/(2 + 1/\tau)$, the pair (τ, s) is **not** in $JS(d)$.*

Proof. If $s > 1/2$ then the result follows from Remark 2.2. So we can assume that $1/(2 + 1/\tau) < s \leq 1/2$. Let us choose some number $a \in (1/(2 + 1/\tau), s)$ and then let $E_- = \{(x, t) : x \in [0, 1]^{d-1}, 0 \leq t \leq a\}$ and $E_+ = \{(x, t) : x \in [0, 1]^{d-1}, 1 - a \leq t \leq 1\}$. Since $a < 1/2$, these two sets are disjoint. Furthermore

$$\lambda([0, 1]^d \setminus E_+ \setminus E_-) = 1 - 2a = \frac{1 - 2a}{a} \min \{ \lambda(E_+), \lambda(E_-) \}$$

or, equivalently,

$$\min \{ \lambda(E_+), \lambda(E_-) \} = \frac{1}{\frac{1}{a} - 2} \lambda([0, 1]^d \setminus E_+ \setminus E_-).$$

Since $a > 1/(2 + 1/\tau)$ it follows that $\frac{1}{\frac{1}{a} - 2} > \tau$ and so

$$(4.22) \qquad \min \{ \lambda(E_+), \lambda(E_-) \} > \tau \lambda([0, 1]^d \setminus E_+ \setminus E_-).$$

We will complete the proof of this lemma by showing that, although the two disjoint measurable subsets E_+ and E_- of $[0, 1]^d$ satisfy (4.22), there does not exist any subcube

W of $[0, 1]^d$ which satisfies $\min \{\lambda(E_+ \cap W), \lambda(E_- \cap W)\} \geq s\lambda(W)$. This will follow from the inequality

$$(4.23) \quad \frac{\min \{\lambda(E_+ \cap W), \lambda(E_- \cap W)\}}{\lambda(W)} \leq a$$

which will be seen to hold for every subcube W of $[0, 1]^d$. This inequality seems intuitively quite obvious, but let us nevertheless give a detailed (and perhaps not optimally elegant) proof.

Let W be a subcube of $[0, 1]^d$ and let θ be its side length. Of course $\theta \in (0, 1]$ and W must be of the form $W = \{(x, t) : x \in W_0, t \in [\beta, \beta + \theta]\}$ where W_0 is some subcube of $[0, 1]^{d-1}$ of side length θ and $[\beta, \beta + \theta] \subset [0, 1]$. We only need to consider the case where $[\beta, \beta + \theta]$ has a non-empty intersection with both of the intervals $[0, a]$ and $[1 - a, 1]$, since otherwise at least one of the two sets $E_+ \cap W$ and $E_- \cap W$ is empty and (4.23) is a triviality. The non-emptiness of the above mentioned two intersections implies that

$$\beta \leq a \text{ and } \beta + \theta \geq 1 - a.$$

When θ and β satisfy all the above mentioned conditions we have that $\lambda(E_- \cap W) = \theta^{d-1}(a - \beta)$ and $\lambda(E_+ \cap W) = \theta^{d-1}(\beta + \theta - (1 - a))$. Therefore

$$(4.24) \quad \min \{\lambda(E_+ \cap W), \lambda(E_- \cap W)\} \leq \theta^{d-1} \sup_{\beta \in \mathbb{R}} [\min \{a - \beta, \beta + \theta - (1 - a)\}].$$

For each fixed choice of θ , the expression in the square brackets on the right side of (4.24) is the minimum of a strictly decreasing function of β and a strictly increasing function of β . Therefore its supremum and thus its maximum is attained when β takes the unique value for which these two functions are equal, namely when $\beta = (1 - \theta)/2$. This shows that

$$(4.25) \quad \frac{\min \{\lambda(E_+ \cap W), \lambda(E_- \cap W)\}}{\lambda(W)} \leq \frac{\theta^{d-1}}{\theta^d} \left(a - \frac{1 - \theta}{2} \right) = \frac{\theta^2}{2} + \left(a - \frac{1}{2} \right) \theta.$$

Finally, the fact that $0 < \theta \leq 1$ implies that the right side of (4.25) is bounded above by

$$\frac{\theta}{2} + \left(a - \frac{1}{2} \right) \theta = a\theta \leq a$$

which establishes (4.23) and so completes the proof of the lemma. $\square$

Remark 4.7. In particular, when $\tau = 1/2$, Lemma 4.6 shows that $(\frac{1}{2}, \frac{1}{4} + \varepsilon) \notin JS(d)$ for every $d \in \mathbb{N}$ and every $\varepsilon > 0$. Thus the second of the three results obtained recently by Ron Holzman (see Remark 1.2), namely that $(\frac{1}{2}, \frac{1}{4}) \in JS(1)$, is best possible, and the first of his three results can also be considered, in some sense, to be best possible. (See also Remark 4.10.) These results tempt one to wonder whether perhaps the property $(\frac{1}{2}, \frac{1}{4}) \in JS(d)$ might hold for *all* $d \in \mathbb{N}$. If so, that would of course also answer Question A(1/2) affirmatively, and in an optimally strong, even “dramatically strong” way. However, the following example, which was suggested by the author and analyzed by Holzman, shows that this is an “impossible dream”. This property fails to hold already for $d = 2$. Therefore (cf. Theorem 4.13 below) it also does not hold for any other $d > 1$.

Let F_+ be the rectangle $F_+ = [0, 1] \times [2/3, 1]$ and let F_- be the union of the two squares $[0, 1/3] \times [0, 1/3]$ and $[2/3, 1] \times [0, 1/3]$. Then (F_+, F_-) is a tame couple in $[0, 1]^2$. The areas of the three disjoint sets F_+ , F_- and $[0, 1]^2 \setminus F_+ \setminus F_-$ are respectively $1/3$, $2/9$ and $4/9$, and this ensures that F_+ and F_- satisfy (4.1) for $\tau = 1/2$ and $d = 2$. (In this example λ will of course always denote two-dimensional Lebesgue measure.) Let $\mathcal{W}$ denote the collection of all closed squares with sides parallel to the axes which are contained in $[0, 1]^2$. For each $W \in \mathcal{W}$ let

$$f(W) = \frac{\min \{\lambda(W \cap F_+), \lambda(W \cap F_-)\}}{\lambda(W)}.$$

The third of Ron Holzman's results is that, for these choices of F_+ and F_- ,

$$\sup \{f(W) : W \in \mathcal{W}\} = \max \{f(W) : W \in \mathcal{W}\} = \sqrt{5} - 2.$$

In view of part (***) of Theorem 4.3, this shows that, indeed, $(1/2, 1/4) \notin JS(2)$ and, furthermore, that $(1/2, s) \notin JS(d)$ for every $s > \sqrt{5} - 2$ and $d \geq 2$.

It is tempting to wonder whether a sequence of appropriate higher dimensional variants of this example, where these dimensions tend to ∞ , might lead to a negative answer to Question A(1/2). Initial attempts to find such a sequence have not yielded anything decisive.

4.3. Some further properties of the set $JS(d)$.

Fact 4.8. *For each $d \in \mathbb{N}$ and each $\tau > 0$ there exists some $s > 0$ such that $(\tau, s) \in JS(d)$.*

Proof. This is a consequence of Theorem 7.8 of [1, pp. 28–29] a.k.a. Theorem 7.7 of [3, pp. 152–153] which, for each $\tau > 0$, provides a positive number s depending on τ and d such that $(\tau, s) \in JS(d)$. (The explicit formula for s will be recalled and used in Theorem 4.11 below. $\square$)

The preceding result ensures that the supremum in the following definition is taken over a non-empty set.

Definition 4.9. For each $d \in \mathbb{N}$ and $\tau > 0$ let

$$\sigma(\tau, d) := \sup \{s > 0 : (\tau, s) \in JS(d)\}.$$

Remark 4.10. Thus the second and third results of Ron Holzman mentioned in Remark 1.2 (cf. also Remark 4.7) can be written as

$$\sigma\left(\frac{1}{2}, 1\right) = \frac{1}{4} \quad \text{and} \quad \sigma\left(\frac{1}{2}, 2\right) \leq \sqrt{5} - 2.$$

We can now readily establish several properties of the set $JS(d)$ and the function $\sigma(\tau, d)$.

Theorem 4.11. *For each fixed $d \in \mathbb{N}$,*

- (i) $(\tau, \sigma(\tau, d)) \in JS(d)$ for each $\tau > 0$.
- (ii) $JS(d) = \{(x, y) : x > 0, 0 < y \leq \sigma(x, d)\}$
- (iii) *The function $x \mapsto \sigma(x, d)$ is non-decreasing and continuous and satisfies*

$$(4.26) \quad \varphi(x, d) \leq \sigma(x, d) \leq \frac{1}{2 + \frac{1}{x}} \text{ for all } x > 0,$$

where

$$\varphi(x, d) = \begin{cases} 2^{-d}(x - x^2)/(1 + x) & , \quad 0 < x \leq \sqrt{2} - 1 \\ 2^{-d}(3 - 2\sqrt{2}) & , \quad x \geq \sqrt{2} - 1. \end{cases}$$

Proof. For part (i) let us fix an arbitrary $\tau > 0$ and first note the obvious fact that

$$(4.27) \quad (\tau, s) \in JS(d) \Rightarrow (\tau, s') \in JS(d) \text{ for all } s' \in (0, s),$$

which implies that $(\tau, s') \in JS(d)$ for every $s' \in (0, \sigma(\tau, d))$. Therefore the sequences $\{\tau_n\}_{n \in \mathbb{N}}$ and $\{s_n\}_{n \in \mathbb{N}}$ which we define by $\tau_n := \tau$ and $s_n := (1 - 2^{-n})\sigma(\tau, d)$ must satisfy $(\tau_n, s_n) \in JS(d)$ for every $n \in \mathbb{N}$. For the proof of part (i) we now simply apply Theorem 4.3 together with part (a) of Lemma 4.5 to the sequences $\{\tau_n\}_{n \in \mathbb{N}}$ and $\{s_n\}_{n \in \mathbb{N}}$. Part (ii) then follows immediately from part (i) and (4.27). For part (iii) we first use another obvious fact, namely that

$$(\tau, s) \in JS(d) \Rightarrow (\tau', s) \in JS(d) \text{ for all } \tau' > \tau$$

to immediately imply that the function $x \mapsto \sigma(x, d)$ is non-decreasing. This latter property means that, in order to show that this function is continuous, it will suffice to show that

$$(4.28) \quad \lim_{n \rightarrow \infty} \sigma((1 + 1/n)\tau, d) \leq \sigma(\tau, d) \text{ and } \lim_{n \rightarrow \infty} \sigma((1 - 1/n)\tau, d) \geq \sigma(\tau, d) \text{ for each } \tau > 0.$$

We obtain the first of these inequalities by again using Theorem 4.3 together with part (a) of Lemma 4.5 to show that the limit of the sequence $\{((1 + 1/n)\tau, \sigma((1 + 1/n)\tau, d))\}_{n \in \mathbb{N}}$ of points in $JS(d)$ must also be a point in $JS(d)$. We next remark that, since $(\tau, \sigma(\tau, d)) \in JS(d)$, it follows from Theorem 3.1, that $((1 - 1/n)\tau, \sigma(\tau, d) - (1 - \sigma(\tau, d))/n) \in JS(d)$ for all sufficiently large $n \in \mathbb{N}$. Therefore $\sigma((1 - 1/n)\tau, d) \geq \sigma(\tau, d) - (1 - \sigma(\tau, d))/n$ for these same values of n . This suffices to prove the second inequality in (4.28) and complete the proof of continuity.

The estimate from below in (4.26) is obtained by once more appealing to Theorem 7.8 of [1, pp. 28–29] a.k.a. Theorem 7.7 of [3, pp. 152–153], and using the fact that, in the terminology introduced just before the statement of that theorem, the collection of sets $\mathcal{Q}(\mathbb{R}^d)$ is 2^d -decomposable. The estimate from above in (4.26) follows from Lemma 4.6.

This completes the proof of part (iii) and therefore of the whole theorem. $\square$

Remark 4.12. We have not bothered to explicitly state another different kind of lower bound for $\sigma(\tau, d)$, in terms of values of $\sigma(\tau', d)$ for appropriate numbers τ' greater than τ , which can be obtained from Theorem 3.1.

The following result seems intuitively completely obvious. But we shall provide an explicit proof.

Theorem 4.13. *The inclusion $JS(d+1) \subset JS(d)$ and consequently also the inequality $\sigma(\tau, d+1) \leq \sigma(\tau, d)$ both hold for every $d \in \mathbb{N}$ and $\tau > 0$.*

Proof. This is one place in this paper where we need to use the more explicit notation λ_d instead of λ to denote d -dimensional Lebesgue measure. We will use Theorem 2.4. (It would perhaps be slightly simpler to use the fact, which is easily explained in [1]

and [3], that the statement of Theorem 2.4 remains true when the word “*d-multi-cubes*” is replaced by “*Lebesgue measurable subsets of $[0, 1]^d$* ”.)

Suppose that $(\tau, s) \in JS(d+1)$. Let F_+ and F_- be two *arbitrarily chosen* disjoint d -multi-cubes which satisfy (2.3), i.e., the inequality

$$(4.29) \quad \min \{ \lambda_d(F_+), \lambda_d(F_-) \} > \tau \lambda_d([0, 1]^d \setminus F_+ \setminus F_-),$$

for this given value of τ . We define two subsets H_+ and H_- of $[0, 1]^{d+1}$ as the cartesian products $H_+ = F_+ \times [0, 1]$ and $H_- = F_- \times [0, 1]$. They are disjoint, since F_+ and F_- are disjoint. For each dyadic subcube E of $[0, 1]^d$, the cartesian product $E \times [0, 1]$ is the union of 2^k dyadic subcubes of $[0, 1]^{d+1}$, where k is such that the side length of E is 2^{-k} . It follows that H_+ and H_- are both $(d+1)$ -multi-cubes.

Now, and also again later, we shall use the very standard facts that

(i) for each bounded closed interval $[a, b]$, the set $E \times [a, b]$ is a Lebesgue measurable subset of $\mathbb{R}^{d+1}$ whenever E is a Lebesgue measurable subset of $\mathbb{R}^d$, and that

(ii) $\lambda_{d+1}(E \times [a, b]) = (b - a)\lambda_d(E)$ for each such E .

These two facts together with the fact that $[0, 1]^{d+1} \setminus H_+ \setminus H_- = ([0, 1]^d \setminus F_+ \setminus F_-) \times [0, 1]$, and together with our assumption that F_+ and F_- satisfy (4.29), imply that

$$\min \{ \lambda_{d+1}(H_+), \lambda_{d+1}(H_-) \} > \tau \lambda_{d+1}([0, 1]^{d+1} \setminus H_+ \setminus H_-).$$

Consequently, by Theorem 2.4, our assumption that $(\tau, s) \in JS(d+1)$ guarantees the existence of a subcube V of $[0, 1]^{d+1}$ for which

$$(4.30) \quad \min \{ \lambda_{d+1}(V \cap H_+), \lambda_{d+1}(V \cap H_-) \} \geq s \lambda_{d+1}(V).$$

We can write V as the cartesian product $V = W \times [a, b]$, where W is a subcube of $[0, 1]^d$ and $[a, b]$ is a subinterval of $[0, 1]$ whose length $b - a$ of course equals the side length of V and of W . Obviously $V \cap H_+ = (W \cap F_+) \times [a, b]$ and $V \cap H_- = (W \cap F_-) \times [a, b]$. So we can again apply the standard facts (i) and (ii), which were recalled in an earlier step of this proof, to obtain the formulae $\lambda_{d+1}(E) = (b - a)\lambda_d(E)$ in the three cases where E is $W \cap F_+$ or $W \cap F_-$ or W . When we substitute these formulae in (4.30) and divide both sides of the inequality by $b - a$, we obtain that the subcube W of $[0, 1]^d$ satisfies (2.4) of Theorem 2.4, i.e., that

$$\min \{ \lambda_d(W \cap F_+), \lambda_d(W \cap F_-) \} \geq s \lambda_d(W).$$

Since F_+ and F_- were chosen arbitrarily, we can once more apply Theorem 2.4 to deduce that $(\tau, s) \in JS(d)$, and so complete the proof of the present theorem. $\square$

5. SOME COMMENTS AND MINOR CORRECTIONS FOR THE PAPERS [1, 3].

In the first paragraph of the proof of Theorem 7.8 [1, p. 30] a.k.a. Theorem 7.7 [3, pp. 154–155] it is shown that it suffices to consider the case where the two sets E_+ and E_- are both compact. The justification of this is a little clearer if in the formula on the third line of the proof one replaces G by $Q \setminus H_+ \setminus H_-$, which is obviously permissible.

We refer to the remarks made above in the course of the proof of Theorem 2.4 for some other small corrections and clarifications of some small issues in [1].

REFERENCES

- [1] M. Cwikel, Y. Sagher and P. Shvartsman, A new look at the John-Nirenberg and John-Strömberg theorems for BMO. Lecture Notes. [arXiv:1011.0766v1](#) [[math.FA](#)] (*Posted on 2 Nov 2010*).
- [2] M. Cwikel, Y. Sagher and P. Shvartsman, A geometrical/combinatorial question with implications for the John-Nirenberg inequality for BMO functions, *Banach Center Publ.* **95** (2011), 45–53 .
- [3] M. Cwikel, Y. Sagher and P. Shvartsman, A new look at the John-Nirenberg and John-Strömberg theorems for BMO, *J. Funct. Anal.* **263** (2012), 129–166.
- [4] R. Holzman, Private Communication.
- [5] F. John and L. Nirenberg, On functions of bounded mean oscillation, *Comm. Pure Appl. Math.* **14** (1961), 415–426.
- [6] A. K. Lerner, The John-Nirenberg inequality with sharp constants, *C. R. Math. Acad. Sci. Paris* **351** (2013), 463–466.
- [7] L. Slavin, The John–Nirenberg constant of BMO^p , $1 \leq p \leq 2$. [arXiv:1506.04969v1](#) [[math.CA](#)] (*Posted on 16 Jun 2015*).
- [8] L Slavin and V. Vasyunin, Inequalities for BMO on α -Trees. *IMRN* (*On line as of 1 Oct 2015*). <http://imrn.oxfordjournals.org/content/early/2015/09/30/imrn.rnv258.full.pdf?keytype=ref&ijkey=tvZXuAfzcBuzC>
- [9] L. Slavin and V. Vasyunin, The John–Nirenberg constant of BMO^p , $p > 2$. [arXiv:1601.03848](#) [[math.CA](#)] (*Posted on 15 Jan 2016*).
- [10] R. Wheeden and A. Zygmund, *Measure and integral. An introduction to real analysis*. Pure and Applied Mathematics, Vol. 43. Marcel Dekker, Inc., New York-Basel, 1977.

DEPARTMENT OF MATHEMATICS, TECHNION - ISRAEL INSTITUTE OF TECHNOLOGY, HAIFA 32000,
ISRAEL

E-mail address: mcwikel@math.technion.ac.il